

MTA 2019

Notation

\mathbb{Z} = the set of integers

$\mathbb{N} = \{n \in \mathbb{Z} : n \geq 1\}$

\mathbb{R} = the set of real numbers

\mathbb{Q} = the set of rational numbers

\mathbb{C} = the set of complex numbers

- (1) (a) Let $\Delta = \{(t_0, t_1, t_2) \in \mathbb{R}^3 : t_0 + t_1 + t_2 = 1 \text{ and } t_i \geq 0 \text{ for } i = 0, 1, 2\}$. Prove that the function $f : [0, 1] \times [0, 1] \rightarrow \Delta$ defined by

$$f(x_1, x_2) = \begin{cases} (x_1, x_2 - x_1, 1 - x_2) & \text{if } x_1 \leq x_2 \\ (x_2, x_1 - x_2, 1 - x_1) & \text{if } x_2 \leq x_1 \end{cases}$$

is continuous.

- (b) Prove that $f(A \times B)$ is closed if A and B are closed subsets of $[0, 1]$.

- (2) Show that

$$\lim_{n \rightarrow \infty} \int_{\alpha}^{\infty} \sqrt{n} e^{-nx^2} dx = \int_{\alpha}^{\infty} \lim_{n \rightarrow \infty} \sqrt{n} e^{-nx^2} dx$$

for $\alpha > 0$ but not for $\alpha = 0$.

- (3) Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ and $f : \mathbb{D} \rightarrow \mathbb{D}$ be a holomorphic function such that f has a zero of order $N \geq 1$ at the origin. Show that $|f(z)| \leq |z|^N$ for all $z \in \mathbb{D}$.

- (4) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a C^∞ -function such that $f(x) = 0$ if and only if $x \in \mathbb{Z}$. Suppose the function $x : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $x'(t) = f(x(t))$ for all $t \in \mathbb{R}$.
- (a) If $\mathbb{Z} \cap \{x(t) : t \in \mathbb{R}\}$ is non-empty, then show that x is constant.
- (b) If $\mathbb{Z} \cap \{x(t) : t \in \mathbb{R}\}$ is the empty set, then show that $\lim_{t \rightarrow \infty} x(t)$ exists and is an integer.

(5) (a) Let X be a Banach space, $x_0 \in X$ and $\varphi_0 \in X^*$. Define $T : X^* \rightarrow X^*$ by $T(\psi) = \psi(x_0)\varphi_0$ for $\psi \in X^*$. Prove that T is compact.

(b) Using part (a) or otherwise, prove that given a two-variable polynomial function a , the operator $A : L^\infty([0, 1], m) \rightarrow L^\infty([0, 1], m)$ (where m denotes the Lebesgue measure) defined by

$$Af(x) = \int_0^1 a(x, y)f(y)dy$$

is compact.

(6) Let $f : \mathbb{R} \rightarrow [0, 1]$ be a continuously differentiable function satisfying $f^2(x) + (f'(x))^2 \geq 1$ for all $x \in \mathbb{R}$ and $f(0) = 1$. Prove that there exists $t > 0$ such that $f'(t) = 0$.

Hint: Note that f attains its maximum at 0.

(7) Let $\{x_n\}_{n \geq 1}$ be a sequence in \mathbb{R} and $\{a_n\}_{n \geq 1}$ be a sequence of positive real numbers satisfying $a_n \uparrow \infty$ as $n \rightarrow \infty$. Further, suppose $\sum_{n=1}^{\infty} \frac{x_n}{a_n}$

converges. Then, show that $\frac{1}{a_n} \sum_{k=1}^n x_k \rightarrow 0$ as $n \rightarrow \infty$.

(8) If f is an entire function such that $\iint_{\mathbb{R}^2} |f(x + iy)| dx dy < \infty$, then prove that $f \equiv 0$.

(9) Let $f : [0, 1] \rightarrow \mathbb{R}$ be a monotonically increasing function (not necessarily continuous) with $f(0) = 0$ and $f(1) = 1$. Suppose μ denotes the Borel measure on $[0, 1]$ such that $\mu((a, b))$ is the cardinality of the set

$$\left\{ x \in [0, 1] \mid a < \lim_{h \rightarrow 0^+} [f(x+h) - f(x)] \leq b \right\} \text{ for all } 0 \leq a < b \leq 1.$$

Prove that $\int_0^1 t^p d\mu < \infty$ for all $p > 1$.

Hint: Consider $\int_0^1 t^{p-1} \mu([t, 1]) dt$.

(10) Let \mathcal{H} be a Hilbert space and \mathcal{K} be a closed subspace of \mathcal{H} . Given any bounded linear functional $f : \mathcal{K} \rightarrow \mathbb{C}$, prove that there exists a **unique** extension $\tilde{f} : \mathcal{H} \rightarrow \mathbb{C}$ of f as a bounded linear functional satisfying $\|\tilde{f}\| = \|f\|$.