

1. (a) Find the limit of the sequence  $\{x_n\}$  as  $n \rightarrow \infty$ , where

$$x_n = \sum_{k=1}^n \frac{1}{\sqrt{n^2 + k}}.$$

- (b) Let  $g$  be a non-vanishing continuous real-valued function on  $[0, \infty)$  such that  $g(x+y) = g(x)g(y)$  for all  $x, y \geq 0$ . Prove that there exists a real number  $a$  such that  $g(x) \equiv e^{ax}$ .

[5+7]=12

2. (a) Let  $f$  be a real-valued function defined as follows:

$$f(x, y) = \begin{cases} \frac{e^{x^2 y^2} - 1}{x^2 + y^2}, & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

Is  $f$  continuous everywhere? Justify your answer.

- (b) Suppose that  $f : \mathbb{R} \mapsto \mathbb{R}$  is a differentiable function such that  $f(x) > f(0)$  and  $f(x) > f(1)$  for some  $x \in (0, 1)$ . Prove that there exists a point  $x_0 \in (0, 1)$  such that  $f'(x_0) = 0$ .

[6+6]=12

3. Let  $P$  be an  $n \times n$  non-singular matrix such that  $I + P + P^2 + \dots + P^n$  is a null matrix. Find the inverse of  $P$  and the eigenvalues of  $P$ .

[6+6]=12

4. (a) Suppose that there are five pairs of shoes in a closet and four shoes are taken out at random. What is the probability that, among the four which are taken out, there is at least one complete pair?

- (b) Two identical independent components having lifetime  $T_1$  and  $T_2$ , respectively, are connected in a parallel system. Suppose that the distributions of both  $T_1$  and  $T_2$  are exponential initially with mean  $1/\lambda$ . But, whenever one component fails, the lifetime distribution of the remaining component changes to exponential with mean  $1/\alpha$ . If  $T$  denotes the overall lifetime of the system, find  $P(T \geq t)$  for any  $t > 0$ .

[5+7]=12

5. (a) Suppose that  $(X, Y)$  has a joint distribution with  $E(Y|X = x) = x^3$  for all  $x \in \mathbb{R}$ . If the marginal distribution of  $X$  is  $\mathcal{N}(0, 1)$ , then prove that

$$\text{Correlation}(X, Y) > 0.$$

(b) Suppose that  $(X, Y) \sim BN(0, 0, 1, 1, \rho)$ , and define

$$Z = \frac{X - Y}{X + Y} \sqrt{\frac{1 + \rho}{1 - \rho}}.$$

Show that the distribution of  $Z$  is symmetric, and find  $P(Z < 0 | X + Y < 0)$ .

[5+3+4]=12

6. Let  $X_1, \dots, X_n$  be independent and identically distributed as  $\text{Bin}(m, p)$ , where both  $m$  and  $p$  are unknown.

- Write down the first two method of moments equations and solve them to find the estimators  $\hat{m}$  and  $\hat{p}$ .
- Show that  $\hat{m}$  and  $\hat{p}$  are consistent for  $m$  and  $p$ , respectively.
- Show that  $\hat{m}$  and  $\hat{p}$  are jointly asymptotically normal in the sense that

$$\sqrt{n} \begin{pmatrix} \hat{m} - m \\ \hat{p} - p \end{pmatrix} \Rightarrow \mathcal{N}_2(0, \Sigma),$$

where  $\Sigma$  is a covariance matrix. Find  $\Sigma$ .

[3+3+6]=12

7. Consider the following linear model:

$$\begin{aligned} Y_1 &= \theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5 + \theta_6 + \epsilon_1 \\ Y_2 &= \theta_1 + \theta_2 + \theta_3 - \theta_4 - \theta_5 - \theta_6 + \epsilon_2 \\ Y_3 &= \theta_1 - \theta_2 - \theta_3 + \theta_4 + \theta_5 - \theta_6 + \epsilon_3 \\ Y_4 &= \theta_1 - \theta_2 - \theta_3 - \theta_4 - \theta_5 + \theta_6 + \epsilon_4 \\ Y_5 &= -\theta_1 + \theta_2 - \theta_3 + \theta_4 - \theta_5 + \theta_6 + \epsilon_5 \\ Y_6 &= -\theta_1 + \theta_2 - \theta_3 - \theta_4 + \theta_5 - \theta_6 + \epsilon_6 \end{aligned}$$

where  $\epsilon = (\epsilon_1, \dots, \epsilon_6)'$ , with  $E(\epsilon) = \mathbf{0}$ , and  $V(\epsilon) = \sigma^2 \mathbb{I}$ .

- Find an unbiased estimator of  $\theta_i$  for all  $i = 1, \dots, 6$ .
- Find the Best Linear Unbiased Estimator (BLUE) of  $\theta_1 - 2\theta_2 + \theta_3$ .

[7+5]=12

8. Consider a linear model

$$y_i = \alpha + \beta x_i + \gamma z_i + \epsilon_i, \quad i = 1, 2, \dots, n,$$

where independent observations  $\{y_i\}$  of a response variable  $Y$  are regressed on two regressors  $X$  and  $Z$  and  $\{\epsilon_i\}$  are unobserved independent and identically distributed with  $E(\epsilon_i) = 0$  and  $Var(\epsilon_i) < \infty$  for all  $i$ .

- a) Write down the normal equation for estimating the parameters  $(\alpha, \beta, \gamma)$  by ordinary least square method. Let us denote the resulting estimates by  $(\hat{\alpha}_{OLS}, \hat{\beta}_{OLS}, \hat{\gamma}_{OLS})$ .
- b) Suppose we alternatively fit the model as follows. First we regress  $Y$  on  $X$  by minimizing  $\sum_i (y_i - \alpha_1 - \beta_1 x_i)^2$  with respect to  $(\alpha_1, \beta_1)$  to get their estimates as  $(\hat{\alpha}_1, \hat{\beta}_1)$  and compute the residuals  $e_i = y_i - \hat{\alpha}_1 - \hat{\beta}_1 x_i$  for each  $i$ . In the next step, we regress  $\{e_i\}$  on  $Z$  by minimizing the criterion  $\sum_i (e_i - \alpha_2 - \gamma_2 z_i)^2$  with respect to  $(\alpha_2, \gamma_2)$  to get their estimates as  $(\hat{\alpha}_2, \hat{\gamma}_2)$ . Show that  $\hat{\alpha}_2 = 0$  and  $\hat{\gamma}_2 = \hat{\gamma}_{OLS}$ .

[3+9]=12

9. Let  $X_1, \dots, X_n$  be independent and identically distributed random variables having distribution function  $F_\theta$ . Suppose there exists a positive integer  $m$  such that  $g(X_1, \dots, X_m)$  is unbiased for  $\theta$  and  $E[g(X_1, \dots, X_m)^2] < \infty$ . Prove that if there exists a UMVUE of  $\theta$  for any  $n > m$ , the variance of this UMVUE must converge to zero as  $n \rightarrow \infty$ .

[12]

10. Suppose that a sample of size  $n$  is drawn using SRSWR from a finite population of  $N$  units, where  $N > n$  and  $N \geq 3$ . Let  $\bar{y}$  be the sample mean of the study variables corresponding to the  $n$  selected units. Now, let us assume that one variate value  $y_1$  corresponding to one unit is known and consequently a simple random sample of size  $n$  without replacement are now drawn from the remaining  $(N - 1)$  units; denote the sample mean of the study variables corresponding to these  $n$  selected units by  $\bar{y}_0$ . Consider the following two estimators for the population total as given by

$$t_1 = N\bar{y}, \quad \text{and} \quad t_2 = (N - 1)\bar{y}_0 + y_1.$$

Prove that

- a)  $t_2$  is unbiased for the population total.
- b)  $Var(t_1) \geq Var(t_2)$ .

[5+7]=12