

1. Let  $g : [0, 1] \rightarrow \mathbb{R}$  be a continuous function such that  $g(1) = 0$ . Show that  $\sup_{x \in [0,1]} |x^n g(x)| \rightarrow 0$  as  $n \rightarrow \infty$ . [12]

2. Let  $f : [0, \infty) \rightarrow [0, \infty)$  be an increasing function. For all  $t > 0$ , define  $g(t) = \left( \int_0^t f(x) dx \right) / t$ . Check whether  $g$  is an increasing function. [12]

3. Suppose  $\mathbf{A}$  and  $\mathbf{B}$  are two  $n \times n$  matrices with real entries such that the sum of their ranks is strictly less than  $n$ . Show that there exists a nonzero column vector  $\mathbf{x} \in \mathbb{R}^n$  such that  $\mathbf{Ax} = \mathbf{Bx} = \mathbf{0}$ . [12]

4. In a tennis match between twin brothers Bikas and Bimal, Bikas wins a set with a probability of 0.6. A best-of-five tennis match stops as soon as a player wins three sets. Assume that the outcomes of the sets are independent.

- (a) If the match lasts for five sets, what is the probability that Bikas will win the match?
- (b) If Bimal wins the match, what is the probability that the match lasts for five sets?

[4+8 = 12]

5. Suppose  $X_1, X_2, \dots, X_n$  are independent and identically distributed discrete uniform random variables taking values  $1, 2, 3, \dots, \theta$ , where  $\theta$  is an unknown *positive integer*.

- (a) Define  $\Psi(x) = xe^x - (x - 1)e^{x-1}$  for all  $x \in \mathbb{R}$ . Find  $\mathbb{E}(\Psi(X_1))$ .
- (b) Find the uniformly minimum variance unbiased estimator (UMVUE) of  $e^\theta$ .

[4+8 = 12]

6. If  $X_1, X_2, \dots, X_n$  are independent and identically distributed  $U(0, \theta)$  random variables, show that their geometric mean is a consistent estimator of  $\theta/e$ .

[12]

7. A bivariate random vector  $(X, Y)$  follows uniform distribution on the square region with four vertices  $(\theta, 0), (0, \theta), (-\theta, 0)$  and  $(0, -\theta)$ , where  $\theta > 0$ . Suppose that  $(X_1, Y_1), \dots, (X_n, Y_n)$  are  $n$  independent copies of  $(X, Y)$ .

- (a) Find a real-valued sufficient statistic for  $\theta$ .  
(b) Find the maximum likelihood estimator of  $\theta$ .

[6+6=12]

8. Consider a population  $\{X_1, X_2, \dots, X_{25}\}$  consisting of 25 observations, where each  $X_i$  ( $i = 1, 2, \dots, 25$ ) takes integer values between 0 and 9 (both inclusive). Let the population mean be 5.4. Consider a simple random sample  $\{x_1, x_2, \dots, x_5\}$  of size 5 drawn without replacement from this population. Define  $\bar{x} = (x_1 + \dots + x_5)/5$ .

- (a) Show that  $\text{Var}(\bar{x}) = \sigma^2/6$ , where  $\sigma^2$  is the population variance.  
(b) Show that  $\text{Var}(\bar{x})$  lies between 0.04 and 3.24.

[3+9=12]

9. Construct a test for  $H_0 : \mu = 0$  against  $H_1 : \mu = 1$  based on a single observation from  $N(\mu, 1)$  such that

$$\mathbb{P}(\text{Type I error}) + 2 \mathbb{P}(\text{Type II error})$$

is minimized. Justify your answer.

[12]

10. Consider a linear regression model  $Y_i = \beta_1 X_{1i} + \beta_2 X_{2i} + \epsilon_i$  ( $i = 1, 2, \dots, n$ ) with two non-stochastic covariates  $X_1$  and  $X_2$ , where  $\epsilon_1, \epsilon_2, \dots, \epsilon_n$  are independent and identically distributed  $N(0, \sigma^2)$  variates. Assume that

$$\sum_{i=1}^n Y_i = \sum_{i=1}^n X_{1i} = \sum_{i=1}^n X_{2i} = 0, \quad \sum_{i=1}^n X_{1i}^2 = \sum_{i=1}^n X_{2i}^2 = 1$$

and  $\sum_{i=1}^n X_{1i} X_{2i} = \alpha$  for some  $\alpha \in (-1, 1)$ .

Let  $\hat{\beta}_1$  and  $\hat{\beta}_2$  be the least squares estimates of  $\beta_1$  and  $\beta_2$ , respectively, such that  $\hat{\beta}_1 > \hat{\beta}_2 > 0$ .

- (a) Show that the sample correlation coefficient between  $Y$  and  $X_1$  is larger than that between  $Y$  and  $X_2$ .
- (b) For any  $(\ell_1, \ell_2) \in \mathbb{R}^2$  with  $\ell_1^2 + \ell_2^2 = 1$ , show that the variance of  $\ell_1 \hat{\beta}_1 + \ell_2 \hat{\beta}_2$  cannot be smaller than  $\sigma^2/(1 + |\alpha|)$ .

[6+6 =12]