

## SAMPLE QUESTIONS (MCQ -TYPE)

Note: For each question there are four suggested answers of which only one is correct.

1. Let  $\{f_n(x)\}$  be a sequence of polynomials defined inductively as

$$\begin{aligned}f_1(x) &= (x-2)^2 \\f_{n+1}(x) &= (f_n(x)-2)^2, \quad n \geq 1.\end{aligned}$$

Let  $a_n$  and  $b_n$  respectively denote the constant term and the coefficient of  $x$  in  $f_n(x)$ . Then

- (A)  $a_n = 4, b_n = -4^n$                       (B)  $a_n = 4, b_n = -4n^2$   
(C)  $a_n = 4^{(n-1)!}, b_n = -4^n$                       (D)  $a_n = 4^{(n-1)!}, b_n = -4n^2$ .
2. If  $a, b$  are positive real variables whose sum is a constant  $\lambda$ , then the minimum value of  $\sqrt{(1+1/a)(1+1/b)}$  is

- (A)  $\lambda - 1/\lambda$       (B)  $\lambda + 2/\lambda$       (C)  $\lambda + 1/\lambda$       (D) none of the above.

3. Let  $x$  be a positive real number. Then
- (A)  $x^2 + \pi^2 + x^{2\pi} > x\pi + (\pi + x)x^\pi$
  - (B)  $x^\pi + \pi^x > x^{2\pi} + \pi^{2x}$
  - (C)  $\pi x + (\pi + x)x^\pi > x^2 + \pi^2 + x^{2\pi}$
  - (D) none of the above.
4. Suppose in a competition 11 matches are to be played, each having one of 3 distinct outcomes as possibilities. The number of ways one can predict the outcomes of all 11 matches such that exactly 6 of the predictions turn out to be correct is
- (A)  $\binom{11}{6} \times 2^5$
  - (B)  $\binom{11}{6}$
  - (C)  $3^6$
  - (D) none of the above.
5. A set contains  $2n+1$  elements. The number of subsets of the set which contain at most  $n$  elements is
- (A)  $2^n$
  - (B)  $2^{n+1}$
  - (C)  $2^{n-1}$
  - (D)  $2^{2n}$ .
6. A club with  $x$  members is organized into four committees such that
- (a) each member is in exactly two committees,
  - (b) any two committees have exactly one member in common.
- Then  $x$  has
- (A) exactly two values both between 4 and 8
  - (B) exactly one value and this lies between 4 and 8
  - (C) exactly two values both between 8 and 16
  - (D) exactly one value and this lies between 8 and 16.
7. Let  $X$  be the set  $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ . Define the set  $\mathcal{R}$  by
- $$\mathcal{R} = \{(x, y) \in X \times X : x \text{ and } y \text{ have the same remainder when divided by } 3\}.$$
- Then the number of elements in  $\mathcal{R}$  is
- (A) 40
  - (B) 36
  - (C) 34
  - (D) 33.
8. Let  $A$  be a set of  $n$  elements. The number of ways, we can choose an ordered pair  $(B, C)$ , where  $B, C$  are disjoint subsets of  $A$ , equals
- (A)  $n^2$
  - (B)  $n^3$
  - (C)  $2^n$
  - (D)  $3^n$ .

9. Let  $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ ,  $n$  being a positive integer. The value of

$$\left(1 + \frac{C_0}{C_1}\right) \left(1 + \frac{C_1}{C_2}\right) \dots \left(1 + \frac{C_{n-1}}{C_n}\right)$$

is

- (A)  $\left(\frac{n+1}{n+2}\right)^n$       (B)  $\frac{n^n}{n!}$       (C)  $\left(\frac{n}{n+1}\right)^n$       (D)  $\frac{(n+1)^n}{n!}$ .

10. The value of the infinite product

$$P = \frac{7}{9} \times \frac{26}{28} \times \frac{63}{65} \times \dots \times \frac{n^3 - 1}{n^3 + 1} \times \dots$$

is

- (A) 1      (B)  $\frac{2}{3}$       (C)  $\frac{7}{3}$       (D) none of the above.

11. The number of positive integers which are less than or equal to 1000 and are divisible by none of 17, 19 and 23 equals

- (A) 854      (B) 153      (C) 160      (D) none of the above.

12. Consider the polynomial  $x^5 + ax^4 + bx^3 + cx^2 + dx + 4$  where  $a, b, c, d$  are real numbers. If  $(1 + 2i)$  and  $(3 - 2i)$  are two roots of this polynomial then the value of  $a$  is

- (A)  $-\frac{524}{65}$       (B)  $\frac{524}{65}$       (C)  $-\frac{1}{65}$       (D)  $\frac{1}{65}$ .

13. The number of real roots of the equation

$$2 \cos\left(\frac{x^2 + x}{6}\right) = 2^x + 2^{-x}$$

is

- (A) 0      (B) 1      (C) 2      (D) infinitely many.

14. Consider the following system of equivalences of integers.

$$x \equiv 2 \pmod{15}$$

$$x \equiv 4 \pmod{21}.$$

The number of solutions in  $x$ , where  $1 \leq x \leq 315$ , to the above system of equivalences is

- (A) 0      (B) 1      (C) 2      (D) 3.



21. Let  $\omega$  denote a complex fifth root of unity. Define

$$b_k = \sum_{j=0}^4 j\omega^{-kj},$$

for  $0 \leq k \leq 4$ . Then  $\sum_{k=0}^4 b_k \omega^k$  is equal to

- (A) 5                      (B)  $5\omega$                       (C)  $5(1 + \omega)$                       (D) 0.

22. Let  $a_n = \left(1 - \frac{1}{\sqrt{2}}\right) \cdots \left(1 - \frac{1}{\sqrt{n+1}}\right)$ ,  $n \geq 1$ . Then  $\lim_{n \rightarrow \infty} a_n$

- (A) equals 1              (B) does not exist              (C) equals  $\frac{1}{\sqrt{\pi}}$               (D) equals 0.

23. Let  $X$  be a nonempty set and let  $\mathcal{P}(X)$  denote the collection of all subsets of  $X$ . Define  $f : X \times \mathcal{P}(X) \rightarrow \mathbb{R}$  by

$$f(x, A) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A. \end{cases}$$

Then  $f(x, A \cup B)$  equals

- (A)  $f(x, A) + f(x, B)$   
(B)  $f(x, A) + f(x, B) - 1$   
(C)  $f(x, A) + f(x, B) - f(x, A) \cdot f(x, B)$   
(D)  $f(x, A) + |f(x, A) - f(x, B)|$

24. The series  $\sum_{k=2}^{\infty} \frac{1}{k(k-1)}$  converges to

- (A)  $-1$                       (B) 1                      (C) 0                      (D) does not converge.

25. The limit  $\lim_{x \rightarrow \infty} \left(\frac{3x-1}{3x+1}\right)^{4x}$  equals

- (A) 1                      (B) 0                      (C)  $e^{-8/3}$                       (D)  $e^{4/9}$

26.  $\lim_{n \rightarrow \infty} \frac{1}{n} \left(\frac{n}{n+1} + \frac{n}{n+2} + \cdots + \frac{n}{2n}\right)$  is equal to

- (A)  $\infty$                       (B) 0                      (C)  $\log_e 2$                       (D) 1

27. Let  $\cos^6 \theta = a_6 \cos 6\theta + a_5 \cos 5\theta + a_4 \cos 4\theta + a_3 \cos 3\theta + a_2 \cos 2\theta + a_1 \cos \theta + a_0$ .  
Then  $a_0$  is

- (A) 0                      (B)  $1/32$ .                      (C)  $15/32$ .                      (D)  $10/32$ .

28. In a triangle  $ABC$ ,  $AD$  is the median. If length of  $AB$  is 7, length of  $AC$  is 15 and length of  $BC$  is 10 then length of  $AD$  equals

- (A)  $\sqrt{125}$                       (B)  $69/5$                       (C)  $\sqrt{112}$                       (D)  $\sqrt{864}/5$ .

29. The set  $\{x : \left|x + \frac{1}{x}\right| > 6\}$  equals the set

- (A)  $(0, 3 - 2\sqrt{2}) \cup (3 + 2\sqrt{2}, \infty)$   
 (B)  $(-\infty, -3 - 2\sqrt{2}) \cup (-3 + 2\sqrt{2}, \infty)$   
 (C)  $(-\infty, 3 - 2\sqrt{2}) \cup (3 + 2\sqrt{2}, \infty)$   
 (D)  $(-\infty, -3 - 2\sqrt{2}) \cup (-3 + 2\sqrt{2}, 3 - 2\sqrt{2}) \cup (3 + 2\sqrt{2}, \infty)$

30. Suppose that a function  $f$  defined on  $\mathbb{R}^2$  satisfies the following conditions:

$$\begin{aligned} f(x+t, y) &= f(x, y) + ty, \\ f(x, t+y) &= f(x, y) + tx \text{ and} \\ f(0, 0) &= K, \text{ a constant.} \end{aligned}$$

Then for all  $x, y \in \mathbb{R}$ ,  $f(x, y)$  is equal to

- (A)  $K(x+y)$ .      (B)  $K - xy$ .      (C)  $K + xy$ .      (D) none of the above.

31. Consider the sets defined by the real solutions of the inequalities

$$A = \{(x, y) : x^2 + y^4 \leq 1\} \quad B = \{(x, y) : x^4 + y^6 \leq 1\}.$$

Then

- (A)  $B \subseteq A$   
 (B)  $A \subseteq B$   
 (C) Each of the sets  $A - B$ ,  $B - A$  and  $A \cap B$  is non-empty  
 (D) none of the above.

32. If a square of side  $a$  and an equilateral triangle of side  $b$  are inscribed in a circle then  $a/b$  equals

- (A)  $\sqrt{2/3}$                       (B)  $\sqrt{3/2}$                       (C)  $3/\sqrt{2}$                       (D)  $\sqrt{2}/3$ .

33. If  $f(x)$  is a real valued function such that

$$2f(x) + 3f(-x) = 15 - 4x,$$

for every  $x \in \mathbb{R}$ , then  $f(2)$  is

- (A)  $-15$                       (B)  $22$                       (C)  $11$                       (D)  $0$ .

34. If  $f(x) = \frac{\sqrt{3}\sin x}{2 + \cos x}$ , then the range of  $f(x)$  is

- (A) the interval  $[-1, \sqrt{3}/2]$                       (B) the interval  $[-\sqrt{3}/2, 1]$   
(C) the interval  $[-1, 1]$                       (D) none of the above.

35. If  $f(x) = x^2$  and  $g(x) = x \sin x + \cos x$  then

- (A)  $f$  and  $g$  agree at no points  
(B)  $f$  and  $g$  agree at exactly one point  
(C)  $f$  and  $g$  agree at exactly two points  
(D)  $f$  and  $g$  agree at more than two points.

36. For non-negative integers  $m, n$  define a function as follows

$$f(m, n) = \begin{cases} n + 1 & \text{if } m = 0 \\ f(m - 1, 1) & \text{if } m \neq 0, n = 0 \\ f(m - 1, f(m, n - 1)) & \text{if } m \neq 0, n \neq 0 \end{cases}$$

Then the value of  $f(1, 1)$  is

- (A)  $4$                       (B)  $3$                       (C)  $2$                       (D)  $1$ .

37. Let  $a$  be a nonzero real number. Define

$$f(x) = \begin{vmatrix} x & a & a & a \\ a & x & a & a \\ a & a & x & a \\ a & a & a & x \end{vmatrix}$$

for  $x \in \mathbb{R}$ . Then, the number of distinct real roots of  $f(x) = 0$  is

- (A)  $1$                       (B)  $2$                       (C)  $3$                       (D)  $4$ .

38. A real  $2 \times 2$  matrix  $M$  such that

$$M^2 = \begin{pmatrix} -1 & 0 \\ 0 & -1 - \varepsilon \end{pmatrix}$$

- (A) exists for all  $\varepsilon > 0$
- (B) does not exist for any  $\varepsilon > 0$
- (C) exists for some  $\varepsilon > 0$
- (D) none of the above is true

39. The eigenvalues of the matrix  $X = \begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix}$  are

- (A) 1, 1, 4
- (B) 1, 4, 4
- (C) 0, 1, 4
- (D) 0, 4, 4.

40. Let  $x_1, x_2, x_3, x_4, y_1, y_2, y_3$  and  $y_4$  be fixed real numbers, not all of them equal to zero. Define a  $4 \times 4$  matrix  $\mathbf{A}$  by

$$\mathbf{A} = \begin{pmatrix} x_1^2 + y_1^2 & x_1x_2 + y_1y_2 & x_1x_3 + y_1y_3 & x_1x_4 + y_1y_4 \\ x_2x_1 + y_2y_1 & x_2^2 + y_2^2 & x_2x_3 + y_2y_3 & x_2x_4 + y_2y_4 \\ x_3x_1 + y_3y_1 & x_3x_2 + y_3y_2 & x_3^2 + y_3^2 & x_3x_4 + y_3y_4 \\ x_4x_1 + y_4y_1 & x_4x_2 + y_4y_2 & x_4x_3 + y_4y_3 & x_4^2 + y_4^2 \end{pmatrix}.$$

Then  $\text{rank}(\mathbf{A})$  equals

- (A) 1 or 2.
- (B) 0.
- (C) 4.
- (D) 2 or 3.

41. Let  $k$  and  $n$  be integers greater than 1. Then  $(kn)!$  is not necessarily divisible by

- (A)  $(n!)^k$ .
- (B)  $(k!)^n$ .
- (C)  $n!.k!$ .
- (D)  $2^{kn}$ .

42. Let  $\lambda_1, \lambda_2, \lambda_3$  denote the eigenvalues of the matrix

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos t & \sin t \\ 0 & -\sin t & \cos t \end{pmatrix}.$$

If  $\lambda_1 + \lambda_2 + \lambda_3 = \sqrt{2} + 1$ , then the set of possible values of  $t$ ,  $-\pi \leq t < \pi$ , is

- (A) Empty set      (B)  $\left\{\frac{\pi}{4}\right\}$       (C)  $\left\{-\frac{\pi}{4}, \frac{\pi}{4}\right\}$       (D)  $\left\{-\frac{\pi}{3}, \frac{\pi}{3}\right\}$ .

43. The values of  $\eta$  for which the following system of equations

$$\begin{aligned} x + y + z &= 1 \\ x + 2y + 4z &= \eta \\ x + 4y + 10z &= \eta^2 \end{aligned}$$

has a solution are

- (A)  $\eta = 1, -2$       (B)  $\eta = -1, -2$       (C)  $\eta = 3, -3$       (D)  $\eta = 1, 2$ .

44. Let  $P_1, P_2$  and  $P_3$  denote, respectively, the planes defined by

$$\begin{aligned} a_1x + b_1y + c_1z &= \alpha_1 \\ a_2x + b_2y + c_2z &= \alpha_2 \\ a_3x + b_3y + c_3z &= \alpha_3. \end{aligned}$$

It is given that  $P_1, P_2$  and  $P_3$  intersect exactly at one point when  $\alpha_1 = \alpha_2 = \alpha_3 = 1$ . If now  $\alpha_1 = 2, \alpha_2 = 3$  and  $\alpha_3 = 4$  then the planes

- (A) do not have any common point of intersection  
 (B) intersect at a unique point  
 (C) intersect along a straight line  
 (D) intersect along a plane.

45. Angles between any pair of 4 main diagonals of a cube are

- (A)  $\cos^{-1} 1/\sqrt{3}, \pi - \cos^{-1} 1/\sqrt{3}$       (B)  $\cos^{-1} 1/3, \pi - \cos^{-1} 1/3$   
 (C)  $\pi/2$       (D) none of the above.

46. If the tangent at the point  $P$  with co-ordinates  $(h, k)$  on the curve  $y^2 = 2x^3$  is perpendicular to the straight line  $4x = 3y$ , then

- (A)  $(h, k) = (0, 0)$   
 (B)  $(h, k) = (1/8, -1/16)$   
 (C)  $(h, k) = (0, 0)$  or  $(h, k) = (1/8, -1/16)$   
 (D) no such point  $(h, k)$  exists.

47. Consider the family  $\mathcal{F}$  of curves in the plane given by  $x = cy^2$ , where  $c$  is a real parameter. Let  $\mathcal{G}$  be the family of curves having the following property: every member of  $\mathcal{G}$  intersects each member of  $\mathcal{F}$  orthogonally. Then  $\mathcal{G}$  is given by

- (A)  $xy = k$  (B)  $x^2 + y^2 = k^2$   
 (C)  $y^2 + 2x^2 = k^2$  (D)  $x^2 - y^2 + 2yk = k^2$

48. Suppose the circle with equation  $x^2 + y^2 + 2fx + 2gy + c = 0$  cuts the parabola  $y^2 = 4ax$ , ( $a > 0$ ) at four distinct points. If  $d$  denotes the sum of ordinates of these four points, then the set of possible values of  $d$  is

- (A)  $\{0\}$  (B)  $(-4a, 4a)$  (C)  $(-a, a)$  (D)  $(-\infty, \infty)$ .

49. The polar equation  $r = a \cos \theta$  represents

- (A) a spiral (B) a parabola (C) a circle (D) none of the above.

50. Let

$$V_1 = \frac{7^2 + 8^2 + 15^2 + 23^2}{4} - \left( \frac{7 + 8 + 15 + 23}{4} \right)^2,$$

$$V_2 = \frac{6^2 + 8^2 + 15^2 + 24^2}{4} - \left( \frac{6 + 8 + 15 + 24}{4} \right)^2,$$

$$V_3 = \frac{5^2 + 8^2 + 15^2 + 25^2}{4} - \left( \frac{5 + 8 + 15 + 25}{4} \right)^2.$$

Then

- (A)  $V_3 < V_2 < V_1$  (B)  $V_3 < V_1 < V_2$   
 (C)  $V_1 < V_2 < V_3$  (D)  $V_2 < V_3 < V_1$ .

51. A permutation of  $1, 2, \dots, n$  is chosen at random. Then the probability that the numbers 1 and 2 appear as neighbour equals

(A)  $\frac{1}{n}$                       (B)  $\frac{2}{n}$                       (C)  $\frac{1}{n-1}$                       (D)  $\frac{1}{n-2}$ .

52. Two coins are tossed independently where  $P(\text{head occurs when coin } i \text{ is tossed}) = p_i, i = 1, 2$ . Given that at least one head has occurred, the probability that coins produced different outcomes is

(A)  $\frac{2p_1p_2}{p_1 + p_2 - 2p_1p_2}$     (B)  $\frac{p_1 + p_2 - 2p_1p_2}{p_1 + p_2 - p_1p_2}$     (C)  $\frac{2}{3}$     (D) none of the above.

53. The number of cars ( $X$ ) arriving at a service station per day follows a Poisson distribution with mean 4. The service station can provide service to a maximum of 4 cars per day. Then the expected number of cars that do not get service per day equals

(A) 4                      (B) 0                      (C)  $\sum_{i=0}^{\infty} iP(X = i + 4)$                       (D)  $\sum_{i=4}^{\infty} iP(X = i - 4)$ .

54. If  $0 < x < 1$ , then the sum of the infinite series  $\frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{3}{4}x^4 + \dots$  is

(A)  $\log \frac{1+x}{1-x}$     (B)  $\frac{x}{1-x} + \log(1+x)$   
(C)  $\frac{1}{1-x} + \log(1-x)$     (D)  $\frac{x}{1-x} + \log(1-x)$ .

55. Let  $\{a_n\}$  be a sequence of real numbers. Then  $\lim_{n \rightarrow \infty} a_n$  exists if and only if

- (A)  $\lim_{n \rightarrow \infty} a_{2n}$  and  $\lim_{n \rightarrow \infty} a_{2n+2}$  exists
- (B)  $\lim_{n \rightarrow \infty} a_{2n}$  and  $\lim_{n \rightarrow \infty} a_{2n+1}$  exist
- (C)  $\lim_{n \rightarrow \infty} a_{2n}$ ,  $\lim_{n \rightarrow \infty} a_{2n+1}$  and  $\lim_{n \rightarrow \infty} a_{3n}$  exist
- (D) none of the above.

56. Let  $\{a_n\}$  be a sequence of non-negative real numbers such that the series  $\sum_{n=1}^{\infty} a_n$  is convergent. If  $p$  is a real number such that the series  $\sum \frac{\sqrt{a_n}}{n^p}$  diverges, then

- (A)  $p$  must be strictly less than  $\frac{1}{2}$
- (B)  $p$  must be strictly less than or equal to  $\frac{1}{2}$
- (C)  $p$  must be strictly less than or equal to 1 but can be greater than  $\frac{1}{2}$
- (D)  $p$  must be strictly less than 1 but can be greater than or equal to  $\frac{1}{2}$ .

57. Suppose  $a > 0$ . Consider the sequence

$$a_n = n\{\sqrt[n]{ea} - \sqrt[n]{a}\}, \quad n \geq 1.$$

Then

- (A)  $\lim_{n \rightarrow \infty} a_n$  does not exist
- (B)  $\lim_{n \rightarrow \infty} a_n = e$
- (C)  $\lim_{n \rightarrow \infty} a_n = 0$
- (D) none of the above.

58. Let  $\{a_n\}$ ,  $n \geq 1$ , be a sequence of real numbers satisfying  $|a_n| \leq 1$  for all  $n$ . Define

$$A_n = \frac{1}{n}(a_1 + a_2 + \cdots + a_n),$$

for  $n \geq 1$ . Then  $\lim_{n \rightarrow \infty} \sqrt{n}(A_{n+1} - A_n)$  is equal to

- (A) 0
- (B) -1
- (C) 1
- (D) none of these.

59. In the Taylor expansion of the function  $f(x) = e^{x/2}$  about  $x = 3$ , the coefficient of  $(x - 3)^5$  is

- (A)  $e^{3/2} \frac{1}{5!}$
- (B)  $e^{3/2} \frac{1}{2^5 5!}$
- (C)  $e^{-3/2} \frac{1}{2^5 5!}$
- (D) none of the above.

60. Let  $\sigma$  be the permutation:

$$\begin{array}{cccccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \\ 3 & 5 & 6 & 2 & 4 & 9 & 8 & 7 & 1, & \end{array}$$

$I$  be the identity permutation and  $m$  be the order of  $\sigma$  i.e.

$$m = \min\{\text{positive integers } n : \sigma^n = I\}.$$

Then  $m$  is

- (A) 8                      (B) 12                      (C) 360                      (D) 2520.

61. Let

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}.$$

Then

- (A) there exists a matrix  $C$  such that  $A = BC = CB$   
 (B) there is no matrix  $C$  such that  $A = BC$   
 (C) there exists a matrix  $C$  such that  $A = BC$ , but  $A \neq CB$   
 (D) there is no matrix  $C$  such that  $A = CB$ .

62. If the matrix

$$A = \begin{bmatrix} a & 1 \\ 2 & 3 \end{bmatrix}$$

has 1 as an eigenvalue, then  $\text{trace}(A)$  is

- (A) 4                      (B) 5                      (C) 6                      (D) 7.

63. Let  $\theta = 2\pi/67$ . Now consider the matrix

$$A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}.$$

Then the matrix  $A^{2010}$  is

- (A)  $\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$                       (B)  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$   
 (C)  $\begin{pmatrix} \cos^{30} \theta & \sin^{30} \theta \\ -\sin^{30} \theta & \cos^{30} \theta \end{pmatrix}$                       (D)  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ .

64. Let the position of a particle in three dimensional space at time  $t$  be  $(t, \cos t, \sin t)$ . Then the length of the path traversed by the particle between the times  $t = 0$  and  $t = 2\pi$  is

- (A)  $2\pi$ .      (B)  $2\sqrt{2}\pi$ .      (C)  $\sqrt{2}\pi$       (D) none of the above.

65. Let  $n$  be a positive real number and  $p$  be a positive integer. Which of the following inequalities is true?

- (A)  $n^p > \frac{(n+1)^{p+1} - n^{p+1}}{p+1}$       (B)  $n^p < \frac{(n+1)^{p+1} - n^{p+1}}{p+1}$   
 (C)  $(n+1)^p < \frac{(n+1)^{p+1} - n^{p+1}}{p+1}$       (D) none of the above.

66. The smallest positive number  $K$  for which the inequality

$$|\sin^2 x - \sin^2 y| \leq K|x - y|$$

holds for all  $x$  and  $y$  is

- (A) 2    (B) 1    (C)  $\frac{\pi}{2}$     (D) there is no smallest positive value of  $K$ ; any  $K > 0$  will make the inequality hold.

67. Given two real numbers  $a < b$ , let

$$d(x, [a, b]) = \min\{|x - y| : a \leq y \leq b\} \quad \text{for } -\infty < x < \infty.$$

Then the function

$$f(x) = \frac{d(x, [0, 1])}{d(x, [0, 1]) + d(x, [2, 3])}$$

satisfies

- (A)  $0 \leq f(x) < \frac{1}{2}$  for every  $x$   
 (B)  $0 < f(x) < 1$  for every  $x$   
 (C)  $f(x) = 0$  if  $2 \leq x \leq 3$  and  $f(x) = 1$  if  $0 \leq x \leq 1$   
 (D)  $f(x) = 0$  if  $0 \leq x \leq 1$  and  $f(x) = 1$  if  $2 \leq x \leq 3$ .

68. Let

$$f(x, y) = \begin{cases} e^{-1/(x^2+y^2)} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

Then  $f(x, y)$  is

- (A) not continuous at  $(0, 0)$   
 (B) continuous at  $(0, 0)$  but does not have first order partial derivatives

- (C) continuous at  $(0, 0)$  and has first order partial derivatives, but not differentiable at  $(0, 0)$   
 (D) differentiable at  $(0, 0)$

69. Consider the function

$$f(x) = \begin{cases} \int_0^x \{5 + |1 - y|\} dy & \text{if } x > 2 \\ 5x + 2 & \text{if } x \leq 2 \end{cases}$$

Then

- (A)  $f$  is not continuous at  $x = 2$   
 (B)  $f$  is continuous and differentiable everywhere  
 (C)  $f$  is continuous everywhere but not differentiable at  $x = 1$   
 (D)  $f$  is continuous everywhere but not differentiable at  $x = 2$ .
70. Let  $w = \log(u^2 + v^2)$  where  $u = e^{(x^2+y)}$  and  $v = e^{(x+y^2)}$ . Then

$$\left. \frac{\partial w}{\partial x} \right|_{x=0, y=0}$$

is

- (A) 0                                      (B) 1                                      (C) 2                                      (D) 4

71. Let

$$f(x, y) = \begin{cases} 1, & \text{if } xy = 0, \\ xy, & \text{if } xy \neq 0. \end{cases}$$

Then

- (A)  $f$  is continuous at  $(0, 0)$  and  $\frac{\partial f}{\partial x}(0, 0)$  exists  
 (B)  $f$  is not continuous at  $(0, 0)$  and  $\frac{\partial f}{\partial x}(0, 0)$  exists  
 (C)  $f$  is continuous at  $(0, 0)$  and  $\frac{\partial f}{\partial x}(0, 0)$  does not exist  
 (D)  $f$  is not continuous at  $(0, 0)$  and  $\frac{\partial f}{\partial x}(0, 0)$  does not exist.
72. The map  $f(x) = a_0 \cos |x| + a_1 \sin |x| + a_2 |x|^3$  is differentiable at  $x = 0$  if and only if
- (A)  $a_1 = 0$  and  $a_2 = 0$                                       (B)  $a_0 = 0$  and  $a_1 = 0$   
 (C)  $a_1 = 0$                                       (D)  $a_0, a_1, a_2$  can take any real value.
73.  $f(x)$  is a differentiable function on the real line such that  $\lim_{x \rightarrow \infty} f(x) = 1$  and  $\lim_{x \rightarrow \infty} f'(x) = \alpha$ . Then

- (A)  $\alpha$  must be 0  
 (B)  $\alpha$  need not be 0, but  $|\alpha| < 1$   
 (C)  $\alpha > 1$   
 (D)  $\alpha < -1$ .

74. Let  $f$  and  $g$  be two differentiable functions such that  $f'(x) \leq g'(x)$  for all  $x < 1$  and  $f'(x) \geq g'(x)$  for all  $x > 1$ . Then

- (A) if  $f(1) \geq g(1)$ , then  $f(x) \geq g(x)$  for all  $x$   
 (B) if  $f(1) \leq g(1)$ , then  $f(x) \leq g(x)$  for all  $x$   
 (C)  $f(1) \leq g(1)$   
 (D)  $f(1) \geq g(1)$ .

75. The length of the curve  $x = t^3$ ,  $y = 3t^2$  from  $t = 0$  to  $t = 4$  is

- (A)  $5\sqrt{5} + 1$   
 (B)  $8(5\sqrt{5} + 1)$   
 (C)  $5\sqrt{5} - 1$   
 (D)  $8(5\sqrt{5} - 1)$ .

76. Given that  $\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$ , the value of

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+xy+y^2)} dx dy$$

is

- (A)  $\sqrt{\pi/3}$  (B)  $\pi/\sqrt{3}$  (C)  $\sqrt{2\pi/3}$  (D)  $2\pi/\sqrt{3}$ .

77. Let  $R$  be the triangle in the  $xy$ -plane bounded by the  $x$ -axis, the line  $y = x$ , and the line  $x = 1$ . The value of the double integral

$$\int \int_R \frac{\sin x}{x} dx dy$$

is

- (A)  $1 - \cos 1$  (B)  $\cos 1$  (C)  $\frac{\pi}{2}$  (D)  $\pi$ .

78. The value of

$$\lim_{n \rightarrow \infty} \left[ (n+1) \int_0^1 x^n \ln(1+x) dx \right]$$

is

- (A) 0 (B)  $\ln 2$  (C)  $\ln 3$  (D)  $\infty$ .

79. Let  $g(x, y) = \max\{12 - x, 8 - y\}$ . Then the minimum value of  $g(x, y)$  as  $(x, y)$  varies over the line  $x + y = 10$  is

- (A) 5 (B) 7 (C) 1 (D) 3.

80. Let  $0 < \alpha < \beta < 1$ . Then

$$\sum_{k=1}^{\infty} \int_{1/(k+\beta)}^{1/(k+\alpha)} \frac{1}{1+x} dx$$

is equal to

- (A)  $\log_e \frac{\beta}{\alpha}$       (B)  $\log_e \frac{1+\beta}{1+\alpha}$       (C)  $\log_e \frac{1+\alpha}{1+\beta}$       (D)  $\infty$ .

81. If  $f$  is continuous in  $[0, 1]$  then

$$\lim_{n \rightarrow \infty} \sum_{j=0}^{[n/2]} \frac{1}{n} f\left(\frac{j}{n}\right)$$

(where  $[y]$  is the largest integer less than or equal to  $y$ )

- (A) does not exist  
(B) exists and is equal to  $\frac{1}{2} \int_0^1 f(x) dx$   
(C) exists and is equal to  $\int_0^1 f(x) dx$   
(D) exists and is equal to  $\int_0^{1/2} f(x) dx$ .

82. The volume of the solid, generated by revolving about the horizontal line  $y = 2$  the region bounded by  $y^2 \leq 2x$ ,  $x \leq 8$  and  $y \geq 2$ , is

- (A)  $2\sqrt{2}\pi$       (B)  $28\pi/3$       (C)  $84\pi$       (D) none of the above.

83. If  $\alpha, \beta$  are complex numbers then the maximum value of  $\frac{\alpha\bar{\beta} + \bar{\alpha}\beta}{|\alpha\beta|}$  is

- (A) 2  
(B) 1  
(C) the expression may not always be a real number and hence maximum does not make sense  
(D) none of the above.

84. For positive real numbers  $a_1, a_2, \dots, a_{100}$ , let

$$p = \sum_{i=1}^{100} a_i \quad \text{and} \quad q = \sum_{1 \leq i < j \leq 100} a_i a_j.$$

Then

(A)  $q = \frac{p^2}{2}$       (B)  $q^2 \geq \frac{p^2}{2}$       (C)  $q < \frac{p^2}{2}$       (D) none of the above.

85. The differential equation of all the ellipses centred at the origin is

(A)  $y^2 + x(y')^2 - yy' = 0$       (B)  $xyy'' + x(y')^2 - yy' = 0$   
 (C)  $yy'' + x(y')^2 - xy' = 0$       (D) none of these.

86. The coordinates of a moving point  $P$  satisfy the equations

$$\frac{dx}{dt} = \tan x, \quad \frac{dy}{dt} = -\sin^2 x, \quad t \geq 0.$$

If the curve passes through the point  $(\pi/2, 0)$  when  $t = 0$ , then the equation of the curve in rectangular co-ordinates is

(A)  $y = 1/2 \cos^2 x$       (B)  $y = \sin 2x$   
 (C)  $y = \cos 2x + 1$       (D)  $y = \sin^2 x - 1$ .

87. If  $x(t)$  is a solution of

$$(1 - t^2) dx - tx dt = dt$$

and  $x(0) = 1$ , then  $x(\frac{1}{2})$  is equal to

(A)  $\frac{2}{\sqrt{3}}(\frac{\pi}{6} + 1)$       (B)  $\frac{2}{\sqrt{3}}(\frac{\pi}{6} - 1)$       (C)  $\frac{\pi}{3\sqrt{3}}$       (D)  $\frac{\pi}{\sqrt{3}}$ .

88. Let  $f(x)$  be a given differentiable function. Consider the following differential equation in  $y$

$$f(x) \frac{dy}{dx} = yf'(x) - y^2.$$

The general solution of this equation is given by

(A)  $y = -\frac{x+c}{f(x)}$       (B)  $y^2 = \frac{f(x)}{x+c}$   
 (C)  $y = \frac{f(x)}{x+c}$       (D)  $y = \frac{[f(x)]^2}{x+c}$ .

89. Let  $y(x)$  be a non-trivial solution of the second order linear differential equation

$$\frac{d^2y}{dx^2} + 2c \frac{dy}{dx} + ky = 0,$$

where  $c < 0$ ,  $k > 0$  and  $c^2 > k$ . Then

- (A)  $|y(x)| \rightarrow \infty$  as  $x \rightarrow \infty$
- (B)  $|y(x)| \rightarrow 0$  as  $x \rightarrow \infty$
- (C)  $\lim_{x \rightarrow \pm\infty} |y(x)|$  exists and is finite
- (D) none of the above is true.

90. The differential equation of the system of circles touching the  $y$ -axis at the origin is

- (A)  $x^2 + y^2 - 2xy \frac{dy}{dx} = 0$
- (B)  $x^2 + y^2 + 2xy \frac{dy}{dx} = 0$
- (C)  $x^2 - y^2 - 2xy \frac{dy}{dx} = 0$
- (D)  $x^2 - y^2 + 2xy \frac{dy}{dx} = 0$ .

91. Suppose a solution of the differential equation

$$(xy^3 + x^2y^7) \frac{dy}{dx} = 1,$$

satisfies the initial condition  $y(1/4) = 1$ . Then the value of  $\frac{dy}{dx}$  when  $y = -1$  is

- (A)  $\frac{4}{3}$
- (B)  $-\frac{4}{3}$
- (C)  $\frac{16}{5}$
- (D)  $-\frac{16}{5}$ .

92. Consider the group

$$G = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} : a, b \in \mathbb{R}, a > 0 \right\}$$

with usual matrix multiplication. Let

$$N = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} : b \in \mathbb{R} \right\}.$$

Then,

- (A)  $N$  is not a subgroup of  $G$
- (B)  $N$  is a subgroup of  $G$  but not a normal subgroup
- (C)  $N$  is a normal subgroup and the quotient group  $G/N$  is of finite order
- (D)  $N$  is a normal subgroup and the quotient group is isomorphic to  $\mathbb{R}^+$  (the group of positive reals with multiplication).

93. Let  $G$  be a group with identity element  $e$ . If  $x$  and  $y$  are elements in  $G$  satisfying  $x^5y^3 = x^8y^5 = e$ , then which of the following conditions is true?

- (A)  $x = e, y = e$
- (B)  $x \neq e, y = e$
- (C)  $x = e, y \neq e$
- (D)  $x \neq e, y \neq e$

94. Let  $G$  be the group  $\{\pm 1, \pm i\}$  with multiplication of complex numbers as composition. Let  $H$  be the quotient group  $\mathbb{Z}/4\mathbb{Z}$ . Then the number of nontrivial group homomorphisms from  $H$  to  $G$  is

(A) 4

(B) 1

(C) 2

(D) 3.